

screening on clinical outcomes by center: A CF patient registry study. *J Cyst Fibros* 2020;19(2):316–20.

4

**Role of hyperglycemia in cystic fibrosis pulmonary exacerbations**

L. Merjaneh<sup>1</sup>, A. Sidhaye<sup>2</sup>, P. Vu<sup>3</sup>, S. Heltshel<sup>1,4</sup>, C. Goss<sup>5,6</sup>, P. Flume<sup>7</sup>, A. Kelly<sup>8</sup>, M. Rosenfeld<sup>1,9</sup>. <sup>1</sup>Pediatrics, University of Washington, Seattle, WA; <sup>2</sup>Division of Endocrinology, Diabetes and Metabolism, The Johns Hopkins University School of Medicine, Baltimore, MD; <sup>3</sup>Cystic Fibrosis Foundation Therapeutics Development Network Coordinating Center, Seattle, WA; <sup>4</sup>Seattle Children's Research Institute, Cystic Fibrosis Foundation Therapeutics Development Network Coordinating Center, Seattle, WA; <sup>5</sup>Division of Pulmonary, Critical Care and Sleep Medicine, Department of Medicine, University of Washington, Seattle, WA; <sup>6</sup>Division of Pulmonary and Sleep Medicine, Department of Pediatrics, University of Washington, Seattle, WA; <sup>7</sup>Medicine and Pediatrics, Medical University of South Carolina, Charleston, SC; <sup>8</sup>Division of Endocrinology and Diabetes, Department of Pediatrics, Children's Hospital of Philadelphia, Philadelphia, PA; <sup>9</sup>Department of Epidemiology, University of Washington School of Medicine, Seattle, WA

**Background:** Hyperglycemia could affect treatment response during cystic fibrosis (CF) pulmonary exacerbations (PEX). We aimed to evaluate hyperglycemia prevalence during CF PEX, associations between hyperglycemia and PEX outcomes, and feasibility of continuous glucose monitoring (CGM) during PEX.

**Methods:** This was an ancillary study to the STOP2 study—a multicenter randomized clinical trial evaluating the efficacy and safety of different durations of intravenous (IV) antibiotics for PEX in adults with CF [1]. Random glucose levels measured clinically during the first 7 to 10 days of PEX treatment were captured. A subset of participants at four sites also consented to undergo CGM for 5 days, beginning at IV antibiotic day 1 to 4. Associations between hyperglycemia, defined as random glucose greater than 140 mg/dL, and changes in weight and lung function between baseline (initiation of IV treatment), visit 2 (day 7–10), and visit 3 (14 days after completion of IV treatment) were evaluated using linear regression after adjustment for confounders.

**Results:** Random glucose measurements were available for 182 STOP2 participants (18.6% of 982 randomized in STOP2). Of these participants, 37% had CF-related diabetes (CFRD), 27% were taking insulin, mean ± SD age was 31.6 ± 10.8 years, and mean ± SD baseline percentage predicted forced expiratory volume in 1 second (FEV1pp) was 53.6 ± 22.5. Hyperglycemia was detected in 44% of these participants, and 61% of those with hyperglycemia had a diagnosis of CFRD. Table 1 shows the mean estimated differences in changes in FEV1pp and weight from baseline to visit 2 and visit 3 between those with and without hyperglycemia and taking and not taking insulin. No differences were detected according to hyperglycemia or insulin use at visit 2 or visit 3. Ten participants who were not taking insulin or oral hypoglycemic agents in the 4 weeks before study underwent CGM. Mean ± SD age was 30.1 ± 6.5 years, 70% were female, and mean ± SD baseline FEV1pp was 48.9 ± 21.0 L. Mean time at greater than 140 mg/dL was 24.6 ± 12.5%, and 9/10 participants spent more than 4.5% of the time at more than 140 mg/dL.

**Conclusions:** Hyperglycemia identified using random glucose testing was prevalent during CF PEX in patients with and without CFRD, but we did not

detect an association between hyperglycemia and changes in lung function or weight with treatment. CGM is feasible and may be a useful tool in the inpatient setting.

**Acknowledgements:** This work was supported by the Cystic Fibrosis Foundation [MERJAN17ABO, MERJA19GE0].

**Reference**

[1] Goss CH, Heltshel SL, West NE, Skalland M, Sanders DB, Jain R, et al. A randomized clinical trial of antimicrobial duration for cystic fibrosis pulmonary exacerbation treatment. *Am J Respir Crit Care Med* 2021;204(11):1295–305.

5

**Comparison of meal-time dosing of rapid-acting insulin using carbohydrate counting versus fixed doses using continuous glucose monitoring in patients with cystic fibrosis-related diabetes**

J. Ullal<sup>1</sup>, T. Abdulwahid<sup>2</sup>, A. Paul<sup>3</sup>, A. Kelly<sup>4</sup>, K. Hughan<sup>4</sup>. <sup>1</sup>Medicine, University of Pittsburgh Medical Center, Pittsburgh, PA; <sup>2</sup>Department of Medicine, University of Pittsburgh Medical Center Mercy Hospital, Pittsburgh, PA; <sup>3</sup>Endocrinologist, Virtua Medical Group, Sewell, NJ; <sup>4</sup>Division of Endocrinology and Diabetes, Department of Pediatrics, Children's Hospital of Philadelphia, Philadelphia, PA

**Background:** Optimizing prandial bolus insulin dosing is essential for cystic fibrosis (CF)-related diabetes (CFRD) management but the traditionally high-calorie diet prescribed to maintain ideal body mass index and the demands of close glucose monitoring make this difficult. This study leveraged continuous glucose monitoring (CGM) to compare fixed meal dosing with carbohydrate count-based insulin dosing.

**Methods:** People with CFRD treated with meal-time rapid-acting insulin were recruited. A dietician provided carbohydrate counting education, and insulin carbohydrate ratio was determined according to total daily insulin dose at the study start. Abbott Freestyle Flash Libre continuous glucose monitors were worn for 14 days. Participants used fixed mealtime insulin dosing for the first 7 days (fixed dosing week, FDW) and carbohydrate counting-based insulin dosing for the second 7 days (carbohydrate count week, CCW) using the 'rule of 500' to calculate insulin-to-carbohydrate ratio. We also compared the frequency and duration of hypo- and hyperglycemia of the two insulin delivery methods.

**Results:** Six (3 female, 3 male; aged 65 ± 7.7; 2 F508del homozygous, 2 F508del heterozygous, 2 with unknown mutation status) of the nine recruited participants had complete data (3 dropped out or had CGM malfunction). All subjects were on triple modulator therapy for at least 12 months. Average CFRD duration was 8.6 ± 2.9 years, glycosylated hemoglobin was 7.3 ± 0.5%, and weight was 55 ± 5.4 kg. Insulin to carbohydrate ratio ranged from 1:10 grams to 1:17 grams. Average sensor glucose was not significantly different between FDW (7.27 ± 0.83 mmol/L) and CCW (7.04 ± 0.88 mmol/L) (p = 0.8). There were no differences in time in range (CCW, 984 ± 86 minutes; FDW, 919 ± 110 minutes; p = 0.6) or number of low-glucose events (FDW, 11; CCW, 9; p = 0.6).

**Conclusions:** Few studies have compared the efficacy of carbohydrate counting for dosing mealtime insulin with that of fixed pre-meal insulin doses. Larger studies that use CGM to optimize insulin to carbohydrate-

**Table 1. (abstract 4):**

Mean estimated difference (95% confidence interval) in changes in lung function and weight from baseline to visit 2 and visit 3 according to hyperglycemia and insulin use

	Change in ppFEV1 at visit 2	P value	Change in ppFEV1 at visit 3	P value	Change in weight at visit 2	P value	Change in weight at visit 3	P value
<b>Hyperglycemia vs. no hyperglycemia</b>	1.34 (-1.39, 4.08)	0.37	-1.15 (-4.01, 1.71)	0.43	0.33 (-0.11, 0.78)	0.15	0.34 (-0.31, 0.99)	0.31
<b>On insulin vs. no insulin</b>	0.41 (-2.33, 3.14)	0.77	-1.71 (-4.36, 0.94)	0.2	0.27 (-0.21, 0.75)	0.27	0.05 (-0.70-0.80)	0.89